



Diamondback Terrapin

North Carolina Wildlife Profiles



Diamondback Terrapin *Malaclemys terrapin*

Referred to as the “wind turtle,” some old-timers who live along the Gulf ascribe ominous powers to the diamondback terrapin. Fishermen claim that capturing a terrapin will cause the wind to blow into a squall and bad luck to befall them. In North Carolina, a diamondback terrapin shell was found in an American Indian shaman’s burial site as part of his medicine kit, no doubt a sacred object.

There are seven subspecies or, as some taxonomists claim, seven races of diamondback terrapins in North America. Two subspecies, the Northern and the Carolina, are found in North Carolina.

Description

Often you will hear a diamondback before observing it in the marsh. The terrapin’s powerful jaws make a popping noise as the terrapin eats the periwinkle snails and other mollusks found in the marsh grass. The hind legs are large, and the toes have extensive webs that are useful for its semi-aquatic existence. They are powerful swimmers and are feisty when picked up. Actively struggling, the diamondback is known to bite a toe or finger.

The diamondback exhibits a spotted pattern on the head and along the scutes, or plates. Even experts have difficulty identifying the seven subspecies. The coloration, patterns and shell characteristics vary greatly among individuals in the wild. Many of the captive terrapins were released in North and South Carolina after the collapse of the commercial market and may have diluted the genetics of native populations.

History and Status

Diamondback terrapins were once so abundant in North Carolina that they were considered a nuisance. Fishermen were sometimes unable to haul catches due to the weight of terrapins caught in their nets. But by 1920, the terrapin was a much sought-after gourmet item, costing \$90 per dozen. The market boomed and farms were established (notably in Beaufort, Carteret County) to breed the species in captivity. After World War I the market declined because the wild populations were overharvested.

Though not endangered, the diamondback terrapin today is listed both state and federally as a species of Special Concern. Because of this status within the state, a special permit is required to possess or collect this species. Loss of habitat and mortality from incidental capture in nets and crab pots pose the greatest threats to terrapin populations.


Diamondback terrapins are beautiful turtles.



Range and Distribution

The diamondback terrapin is found along the states of the Atlantic and Gulf coasts from Cape Cod, Massachusetts, to Corpus Christi Bay, Texas. The Northern diamondback terrapin is found from the northern shore of Cape Cod south to Cape Hatteras, North Carolina. From here south to Volusia County, Florida, intergradation with the Carolina diamondback terrapin occurs.

Range Map



Diamondback Terrapin

Wildlife Profiles—North Carolina Wildlife Resources Commission

Habitat and Habits

Diamondback terrapins are adapted for life in brackish and salt water. They typically have high site fidelity, meaning they live in one area throughout their lives, and these areas include protected waters behind barrier islands, salt marshes, estuaries and tidal creeks, and flats hidden among the marsh and cord grass. Natural predators include alligators, sharks, raptors, otters, toad fish, and crabs. At high tide, they swim about the marshes in search of food; at low tide, they are found nearly or entirely buried in the mud or hidden under drift. Diamondback terrapins can hold their breath from 45 minutes (during summer) to five hours.

Sometimes groups of this species are found on exposed mudflats, basking in the sun. Mating occurs in the spring when individuals move from open lagoons into small canals and ditches. The larger female will carry her male counterpart upon her back during the courtship ritual. Hatchlings emerge from their eggs in the autumn and sometimes remain buried in the nest over winter or emerge and bury in the mud into the next spring. Adults also hibernate during the winter, submerged in the mud of tidal creeks in deep water where the mud is less likely to freeze. With the onset of warmer weather, hatchlings seek out and remain in dense cover such as marsh grasses and tidal wrack, presumably until they reach a larger size. Diamondback terrapins are mainly carnivorous and while their favorite food is periwinkle, they also eat other mollusks, crustaceans, insects, annelids and the occasional tender shoots or rootlets.

People Interactions

Diamondback terrapins are beautiful turtles. Because of their special concern status, diamondback terrapins are protected from harvest in North Carolina. Humans affect terrapin populations in very pronounced ways. Terrapins are drowned each year in crab pots, especially derelict traps lost in the water. These death traps remain in the water where terrapins are attracted to the bait or to each other. If an individual enters a crab pot, others will follow and eventually drown. One crab pot, found in North Carolina, contained 29 decomposing terrapins. Adult females are typically too large to enter crab pots so adult males and young females are usually the casualties. Terrapin biologists advocate the use of BRDs (bycatch reduction devices), which prevent smaller terrapins from entering crab pots. The increase in raccoons and other predators associated with human development may increase adult, nest, and hatchling depredation.

Wild Facts

Classification

Class: Reptilia

Order: Testudines

Family: Emydidae

Average Length

Length: male 4 in.–5 in.; female 6 in.–9 in.

Weight: 1.5 lbs. (females) 0.5 lbs. (males)

Food

Crustaceans, mollusks, fish, insects, occasionally tender shoots and rootlets

Breeding

Mating occurs in the water in spring. Eggs are laid in late April–July. Clutch size varies from 4–18 oblong eggs. Eggs laid in chamber dug 6 in.–8 in. below surface; nesting occurs in the daytime during high tides.

Incubation periods are dependent on temperature and vary from 61–104 days for certain nests. Age at sexual maturity is estimated at 5–8 years.

Young

Length just over one in.;

weight 6 g.–10 g.

Life Expectancy


May exceed 40 years.

Diamondback Terrapin

Wildlife Profiles—North Carolina Wildlife Resources Commission

NCWRC Interaction

The diamondback terrapin is not only a species of concern in North Carolina but is also a priority species in the N. C. Wildlife Action Plan. Priority species are targeted for conservation action in the hopes of preventing the diamondback from becoming endangered. Management of this species is a challenge for NCWRC due to its habitat preferences and involves a coordinated effort among multiple agencies and partnerships to provide continued protection from the threats to its survival. These threats include derelict crab pot drowning, pollution, and development pressures.



Q&A—Resources for Teachers

1. What is the largest threat to diamondback terrapins?

Unattended crab pots.

2. What do diamondback terrapins eat?

Mollusks (especially saltmarsh periwinkle), crustaceans, worms, insects, fish, and occasionally tender roots.

3. Which gender, male or female terrapins, are most affected by crab pots and why?

Males, because they are smaller and can more easily enter traps.

References

Pritchard, Peter. *Encyclopedia of Turtles* (T.F.H. Publications, 1979).

Rudloe, Jack. *Time of the Turtle* (E.P. Dutton, 1979).

Buhlmann, Kurt, Tracy Tuberville, and Whit Gibbons. *Turtles of the Southeast*, (University of Georgia Press, 2008)

Palmer, W. and A. Braswell. 1995. *Reptiles of North Carolina*. (University of North Carolina Press, 1995).

Credits

Written by Therese Conant; updated by Kendrick Weeks, Leigh Anne Harden, and Kimberly Burge NCWRC .

Photos by NCWRC and courtesy of Ohio University. Illustrated by J.T. Newman.

Produced by the Division of Conservation Education, N.C. Wildlife Resources Commission.

The Wildlife Resources Commission is an Equal Opportunity Employer and all wildlife programs are administered for the benefit of all North Carolina citizens without prejudice toward age, sex, race, religion or national origin. Violations of this pledge may be reported to the Equal Opportunity Employment Officer, N.C. Wildlife Resources Commission.

1751 Varsity Drive, Raleigh, N.C. 27606. (919) 707-0101.